

Comment on ‘Spin Decoherence in Superconducting Atom Chips’

Stefan Scheel, E. A. Hinds, and P. L. Knight

*Quantum Optics and Laser Science, Blackett Laboratory, Imperial College London,
Prince Consort Road, London SW7 2BW, United Kingdom*

(Dated: February 1, 2008)

We comment on a recent paper [Phys. Rev. Lett. **97**, 070401 (2006)] concerning rubidium atoms trapped near a superconducting niobium surface at ~ 4 K. This seeks to calculate the rate of atomic spin flips induced by thermal magnetic noise. We point out that the calculation is in error by a large factor because it is based on the two-fluid model of superconductivity. This model gives a poor description of electromagnetic dissipation just below the critical temperature because it cannot incorporate the case II coherences of a fuller quantum theory.

PACS numbers: 03.65.Yz, 03.75.Be, 34.50.Dy, 42.50.Ct

A recent publication [1] discusses atoms magnetically trapped near a superconducting surface and attempts to calculate the spin flip rate due to thermal magnetic noise. The Letter recounts the formalism developed in [2, 3], largely verbatim, and then applies it to the two-fluid model of superconductivity. This leads the authors to conclude that Rb atoms near a superconducting niobium surface enjoy a lifetime increase of 10^5 as the temperature is lowered by one or two degrees below $T_c \simeq 8$ K. This disagrees dramatically with previous estimates of the effect [3, 4] that were based on measured electromagnetic dissipation in niobium. Here we point out that the use of the two-fluid model has led to an erroneous result.

In the two-fluid model, the density of the normal component scales as $(T/T_c)^4$. As the temperature drops, this rapidly suppresses all dissipative processes, giving a very large derivative just below T_c . This is precisely the behavior shown in figure 2 of [1]. However, it is well known that the two-fluid model provides a poor description of nuclear relaxation and electromagnetic absorption in the temperature range just below T_c . This is because of quantum interference effects in the relevant low-energy scattering processes, which lead to the so-called case II coherence factors. These cause the dissipation to *increase* as the temperature is lowered below T_c before it declines again in the low temperature limit. This behavior was first observed in the case of nuclear-spin relaxation [5], and is called the Hebel-Slichter peak. Chapter 3 of Tinkham’s book [6] gives a very accessible introduction to this subject. There he points out that “the ability of BCS pairing theory, with its coherence factors, to explain this difference [from the two-fluid model] in a natural way was one of the key triumphs which validated the theory”.

More recently, the same type of behavior has been confirmed experimentally for electromagnetic absorption [7], where the case II coherence factors have been observed in superconducting Nb and Pb over the range $T_c/4 \lesssim T < T_c$. These are not pure BCS superconductors but have to be described by the strong-coupling Eliash-

berg theory. Nevertheless, their temperature-dependent absorption exhibits the same type of peaked behavior, which disagrees with the two-fluid model used in [1].

The predictions for spin flip lifetimes made in previous papers [3, 4] are based on experimental results which explicitly confirm the existence of the case II coherence factor in niobium [7]. The main enhancement of lifetime at 4 K compared with room temperature (~ 100) is due to the smaller number of thermal photons per mode. In addition Ref. [3] estimates that the lifetime should increase by a further factor of ~ 10 as the temperature is lowered from T_c to $T_c/2$, in accordance with BCS theory and with the experimental evidence. By contrast, Ref. [1] predicts an increase of 10^5 , not this factor of 10, when cooling only by one or two degrees below T_c . This is an artifact of the two-fluid model. Although the more realistic calculation gives a smaller enhancement of the lifetime, the effect is nevertheless exceedingly significant for the future of superconducting atom chips [8] as low-decoherence quantum devices.

-
- [1] B.S.K. Skagerstam, U. Hohenester, A. Eiguren, and P.K. Rekdal, Phys. Rev. Lett. **97**, 070401 (2006).
 - [2] P.K. Rekdal, S. Scheel, P.L. Knight, and E.A. Hinds, Phys. Rev. A **70**, 013811 (2004).
 - [3] S. Scheel, P.K. Rekdal, P.L. Knight, and E.A. Hinds, Phys. Rev. A **72**, 042901 (2005).
 - [4] C. Henkel, Eur. Phys. J. D **35**, 59 (2005).
 - [5] L.C. Hebel and C.P. Slichter, Phys. Rev. **107**, 901 (1957); *ibid.* **113**, 1504 (1959).
 - [6] M. Tinkham, *Introduction to Superconductivity*, 2nd edition (McGraw-Hill, New York, 1996).
 - [7] O. Klein, E.J. Nicol, K. Holczer, and G. Grüner, Phys. Rev. B **50**, 6307 (1994).
 - [8] T. Nirrengarten, A. Qarry, C. Roux, A. Emmert, G. Nogues, M. Brune, J.-M. Raimond, and S. Haroche, arXiv:quant-ph/0610019 (2006).